


A DIAGNOSTIC APPROACH FOR PATIENTS WITH SUSPECTED CARDIAC AMYLOIDOSIS THAT INCLUDES TESTING FOR MONOCLONAL PROTEIN FOLLOWED BY SCINTIGRAPHY AND/OR BIOPSY¹



Reprinted with permission from Maurer MS, Bokhari S, Damy T, et al. Expert consensus recommendations for the suspicion and diagnosis of transthyretin cardiac amyloidosis. *Circ Heart Fail.* 2019;12:e006075. doi:10.1161/CIRCHEARTFAILURE.119.006075 © 2019 American Heart Association, Inc. All rights reserved.

- ^{99m}Tc-pyrophosphate (^{99m}Tc-PYP) is a noninvasive radioactive tracer utilized as an adjunct in the diagnosis of ATTR-CM, though not FDA approved for that use[†]
- Both planar and SPECT imaging should be reviewed and interpreted using visual and quantitative approaches¹

*If fat pad is negative, biopsy of involved organ is required.

†Please consult individual labeling for risks.

ATTRm, mutant transthyretin amyloidosis; MGUS, monoclonal gammopathy of undetermined significance; MRI, magnetic resonance imaging; SPECT, single-photon emission computed tomography.

Reference: 1. Maurer MS, Bokhari S, Damy T, et al. Expert consensus recommendations for the suspicion and diagnosis of transthyretin cardiac amyloidosis. *Circ Heart Fail.* 2019;12:e006075. doi:10.1161/CIRCHEARTFAILURE.119.006075

© 2021 Pfizer Inc. All rights reserved. January 2021 PP-VDM-TWN-0241-202302

